

$T = 0$ Partition Functions for Potts Antiferromagnets on Möbius Strips and Effects of Graph Topology

Robert Shrock*

Institute for Theoretical Physics
State University of New York
Stony Brook, N. Y. 11794-3840

Abstract

We present exact calculations of the zero-temperature partition function of the q -state Potts antiferromagnet (equivalently the chromatic polynomial) for Möbius strips, with width $L_y = 2$ or 3 , of regular lattices and homeomorphic expansions thereof. These are compared with the corresponding partition functions for strip graphs with (untwisted) periodic longitudinal boundary conditions.

*email: robert.shrock@sunysb.edu

The chromatic polynomial $P(G, q)$ counts the number of ways that one can color a graph G with q colors such that no two adjacent vertices have the same color [1] (for reviews, see [2]). The minimum number of colors needed for this coloring of G is the chromatic number, $\chi(G)$. Besides its intrinsic mathematical interest, the chromatic polynomial has an important connection with statistical mechanics since it is the zero-temperature partition function of the q -state Potts antiferromagnet (AF) [3, 4] on G : $P(G, q) = Z(G, q, T = 0)_{PAF}$. The Potts AF exhibits nonzero ground-state entropy $S_0 \neq 0$ (without frustration) for sufficiently large q on a given lattice graph and is thus an exception to the third law of thermodynamics (e.g. [5]). This is equivalent to a ground state degeneracy per site $W > 1$, since $S_0 = k_B \ln W$. Denoting the number of vertices of G as $n = v(G)$ and $\{G\} = \lim_{n \rightarrow \infty} G$, we have¹

$$W(\{G\}, q) = \lim_{n \rightarrow \infty} P(G, q)^{1/n} \quad (1)$$

Let us consider strips of regular lattices with variable length and fixed width (with the longitudinal and transverse directions taken to be the x and y directions, respectively.) An obvious and important question concerns the effect of boundary conditions (BC's) on the $T = 0$ partition function. We study this effect here via new calculations of exact $T = 0$ partition functions for arbitrarily long Möbius strips of regular lattice graphs and certain homeomorphic expansions thereof, together with a comparison with the corresponding partition functions for cyclic strips, i.e., strips with (untwisted) periodic longitudinal boundary conditions. These strips all have free transverse boundary conditions (for a comparison of the effects of free versus periodic transverse boundary conditions, see [19]). The different longitudinal boundary conditions also entail different topology: the cyclic strips (topologically $\sim$ cylinder) are oriented, while the Möbius strips are unoriented. We use the symbols FBC_x , $PBC_x = cyc.$, and $TPBC_x = Mb.$ for the free, periodic, cyclic, and Möbius longitudinal boundary conditions, respectively. Some related works on calculations of chromatic polynomials for families of graphs are Refs. [9]-[28]; see also the list in [29].

Since $P(G, q)$ is a polynomial, one can generalize q from $\mathbb{Z}_+$ to $\mathbb{C}$. The zeros of $P(G, q)$ in the complex q plane are called chromatic zeros. Their accumulation set in the limit $n \rightarrow \infty$, denoted $\mathcal{B}$, is the continuous locus of points where $W(\{G\}, q)$ is nonanalytic [13, 16, 6], [18]-[25]. The maximal region in the complex q plane to which one can analytically continue the function $W(\{G\}, q)$ from physical values where there is nonzero ground state entropy is denoted R_1 . The maximal value of q where $\mathcal{B}$ crosses or intersects with the (positive) real axis is denoted $q_c(\{G\})$.

¹ At certain special points q_s (typically $q_s = 0, 1, \dots, \chi(G)$), one has the noncommutativity of limits $\lim_{q \rightarrow q_s} \lim_{n \rightarrow \infty} P(G, q)^{1/n} \neq \lim_{n \rightarrow \infty} \lim_{q \rightarrow q_s} P(G, q)^{1/n}$, and hence it is necessary to specify the order of the limits in the definition of $W(\{G\}, q_s)$ [6]. As in Ref. [6] and our other works, we shall use the first order of limits here; this has the advantage of removing certain isolated discontinuities in W .

We label a particular type of strip graph as G_s and the specific graph of length $L_x = m$ repeated subgraph units, e.g. columns of squares in the case of the square strip and hexagons in the case of the kagomé strip, as $(G_s)_m$. If one thinks of the graph as embedded on a rectangular strip of paper, with its longitudinal ends glued with the same or opposite orientation, L_x is the length of this strip in subgraph units. (Note that, e.g., for the Möbius square strip, if one starts at a vertex on a horizontal boundary edge and traverses a path in the longitudinal direction, the pathlength required to return to the original vertex is $2L_x$.) For m greater than a minimal value², the cyclic and Möbius strips of a given $(G_s)_m$ have the same number of vertices n , edges (bonds) e , girth g (length of minimum closed circuit on G_s) and number k_g of circuits of length g . One has $n = t_s m$ where t_s depends on G_s . Writing

$$P((G_s), q) = \sum_{j=0}^{n-1} (-1)^j h_{n-j} q^{n-j} \quad (2)$$

and using the results that [2, 10] $h_{n-j} = \binom{e}{j}$ for $0 \leq j < g-1$ (whence $h_n = 1$ and $h_{n-1} = e$) and $h_{n-(g-1)} = \binom{e}{g-1} - k_g$, it follows that for m greater than the above-mentioned minimal value, these h_j 's are the same for the cyclic and Möbius strips of each type G_s . For a given G_s , as m increases, the h_{n-j} 's for the cyclic and Möbius strips become equal for larger j (see below).

A generic form for chromatic polynomials for recursively defined families of graphs, of which strip graphs G_s are special cases, is

$$P(G_s, q) = \sum_{j=1}^{N_a} c_j(q) (a_j(q))^m \quad (3)$$

where $c_j(q)$ and $a_j(q)$ depend on the type of strip graph G_s but are independent of m . Note that some c_j and a_j may be nonpolynomial algebraic functions of q , in which case certain cancellations occur to yield the polynomial P [20, 21]. The simplest cyclic strip is the circuit graph with m vertices, C_m , for which $P(C_m, q) = (q-1)^m + (q-1)(-1)^m$. The chromatic polynomials for the cyclic and twisted cyclic ladder graphs were calculated in [9] and have $N_a = 4$. W and $\mathcal{B}$ were given for these and other families in [6] (see also [16]). Various iterative methods have been used in the past to calculate chromatic polynomials for recursive families of graphs [9], [13]-[16], [6], [18]-[27].³ A method that we have found useful to compute $P((G_s)_m, q)$ for a variety of strips with FBC_x [20, 21] and $(\text{T})\text{PBC}_x$ [23, 24, 25]

² For example, for $m \leq 4$ ($m \leq 3$) the $L_y = 2$ cyclic (Möbius) square strips exhibit special behavior regarding g and k_g ; for m larger than these respective values, they both have $g = 4$ and $k_g = m$.

³The method used in [12, 27] can also be used to obtain rigorous bounds on W for 2D lattices [12, 30, 31].

makes use of a generating function

$$\Gamma(G_s, q, x) = \sum_{m=m_0}^{\infty} P((G_s)_m, q) x^{m-m_0} \quad (4)$$

where $\Gamma(G_s, q, x) = \mathcal{N}(G_s, q, x)/\mathcal{D}(G_s, q, x)$ with

$$\mathcal{N}(G_s, q, x) = \sum_{j=0}^{\deg_x \mathcal{N}} A_j(q) x^j \quad (5)$$

and

$$\mathcal{D}(G_s, q, x) = 1 + \sum_{j=1}^{\deg_x \mathcal{D}} b_j(q) x^j = \prod_{j=1}^{\deg_x \mathcal{D}} (1 - \lambda_j(q) x) \quad (6)$$

where m_0 , A_j , and λ_j depend on the type of strip, and A_j and b_j are polynomials in q ($m_0 = 2$ for the square and homeomorphically expanded square strips and $m_0 = 1$ for the kagomé strips considered below). Γ is calculated by means of standard theorems for chromatic polynomials such as the deletion-contraction theorem. The a_j and c_j in (3) can be determined from Γ using eqs. (2.17)-(2.21) of [21] (taking into account value of m_0 for a given strip). For a specified G_s , the λ 's in $\mathcal{D}$ that contribute to P are identical to the⁴ a_j 's. For a given type of strip, we shall study the effect of cyclic versus Möbius longitudinal boundary conditions on (i) the set of a_j , (ii) the W function; (iii) the locus $\mathcal{B}$, (iv) the set of c_j , and (v) χ . Quantities (i)-(iv) depend only on G_s while (v) also depends on m .⁵

We consider first strips of the square (sq) lattice of length $L_x = m \geq 2$ and width L_y vertices. The $L_y = 1, 2$ cases have been mentioned above. For the cyclic $L_y = 3$ strip, P , W , and $\mathcal{B}$ were calculated in [24]. Our new result for the $L_y = 3$ Möbius strip is

$$\begin{aligned} P(sq(L_y = 3, Mb.)_m, q) = & (q^2 - 3q + 1)(-1)^m - (q - 1)^m + (q - 2)^m - (q - 4)^m \\ & - (q - 1)[-(q - 2)^2]^m + [(\lambda_{sq,6})^m + (\lambda_{sq,7})^m] + (q - 1)[(\lambda_{sq,8})^m + (\lambda_{sq,9})^m + (\lambda_{sq,10})^m] \end{aligned} \quad (7)$$

where

$$\lambda_{sq,(6,7)} = \frac{1}{2} \left[(q - 2)(q^2 - 3q + 5) \pm \left\{ (q^2 - 5q + 7)(q^4 - 5q^3 + 11q^2 - 12q + 8) \right\}^{1/2} \right] \quad (8)$$

⁴ For our square lattice strips, all λ_j 's contribute to P . For the kagomé strip and homeomorphically expanded square strips considered here, there are, respectively, three λ 's and one λ that do not contribute to P [24].

⁵ It is also of interest to compare chromatic polynomials for lattice strips with cyclic versus free longitudinal boundary conditions; see [20, 21], [23]-[25].

and $\lambda_{sq,j}$, $j = 8, 9, 10$, are the roots of the cubic equation

$$\xi^3 + b_{sq,21}\xi^2 + b_{sq,22}\xi + b_{sq,23} = 0 \quad (9)$$

with

$$b_{sq,21} = 2q^2 - 9q + 12 \quad (10)$$

$$b_{sq,22} = q^4 - 10q^3 + 36q^2 - 56q + 31 \quad (11)$$

$$b_{sq,23} = -(q-1)(q^4 - 9q^3 + 29q^2 - 40q + 22) . \quad (12)$$

For comparison, the cyclic $L_y = 3$ result is [24]

$$\begin{aligned} P(sq(L_y = 3, cyc.)_m, q) &= (q^3 - 5q^2 + 6q - 1)(-1)^m \\ &+ (q^2 - 3q + 1) \left[(q-1)^m + (q-2)^m + (q-4)^m \right] + (q-1) [-(q-2)^2]^m \\ &+ \left[(\lambda_{sq,6})^m + (\lambda_{sq,7})^m \right] + (q-1) \left[(\lambda_{sq,8})^m + (\lambda_{sq,9})^m + (\lambda_{sq,10})^m \right] . \end{aligned} \quad (13)$$

We discuss these further below.

We next consider strips of the kagomé (kg) lattice comprised of m hexagons with each adjacent pair sharing two triangles as adjacent polygons. For the Möbius kagomé strip with $L_y = 2$ we calculate

$$\begin{aligned} P(kg(L_y = 2, Mb.)_m, q) &= -(q-4)^m - (q-1) \left[(q-1)(q-2)^2 \right]^m \\ &+ \left[(\lambda_{kg,1})^m + (\lambda_{kg,2})^m \right] + (q-1) \left[(\lambda_{kg,3})^m + (\lambda_{kg,4})^m \right] \end{aligned} \quad (14)$$

where

$$\begin{aligned} \lambda_{kg,(1,2)} &= \frac{1}{2} \left[q^4 - 6q^3 + 14q^2 - 16q + 10 \right. \\ &\left. \pm \left\{ q^8 - 12q^7 + 64q^6 - 200q^5 + 404q^4 - 548q^3 + 500q^2 - 292q + 92 \right\}^{1/2} \right] \end{aligned} \quad (15)$$

$$\lambda_{kg,(3,4)} = \frac{1}{2} \left[q^3 - 7q^2 + 19q - 20 \pm \left\{ q^6 - 14q^5 + 83q^4 - 278q^3 + 569q^2 - 680q + 368 \right\}^{1/2} \right] . \quad (16)$$

In contrast, the chromatic polynomial of the regular cyclic $L_y = 2$ kagomé strip is [24]

$$P(kg(L_y = 2, cyc.)_m, q) = (q^2 - 3q + 1)(q-4)^m + (q-1) \left[(q-1)(q-2)^2 \right]^m$$

$$+ \left[(\lambda_{kg,1})^m + (\lambda_{kg,2})^m \right] + (q-1) \left[(\lambda_{kg,3})^m + (\lambda_{kg,4})^m \right]. \quad (17)$$

It is also of interest to study homeomorphic expansions of the cyclic square strip.⁶ We start with the cyclic or Möbius $L_y = 2$ square strip and add $k-2$ vertices to each upper bond (edge). Define

$$D_k = \frac{P(C_k, q)}{q(q-1)} = \sum_{s=0}^{k-2} (-1)^s \binom{k-1}{s} q^{k-2-s} \quad (18)$$

where $P(C_k, q) = (q-1)^k + (q-1)(-1)^k$ is the chromatic polynomial for the circuit graph with k vertices. For the twisted cyclic strip we calculate

$$P((Ch)_{(k,2),Mb.,m}, q) = -(-1)^{km} + (D_{k+2})^m + \left(\frac{D_{k+1} + (-1)^k q}{\lambda_{h,4} - \lambda_{h,5}} \right) \left[(\lambda_{h,4})^m - (\lambda_{h,5})^m \right] \quad (19)$$

where

$$\lambda_{h,(4,5)} = \frac{1}{2} \left[-D_{k+1} - (-1)^k (q-2) \pm \left\{ [D_{k+1} - (-1)^k (q-2)]^2 + 4D_k (-1)^k \right\}^{1/2} \right], \quad (20)$$

h means homeomorphic, and (k_1, k_2) refers to the respective number of vertices (counting the end ones) on the upper and lower horizontal bounds between each vertical bond.

For the cyclic case [24],

$$P((Ch)_{(k,2),cyc.,m}, q) = (q^2 - 3q + 1)(-1)^{km} + (D_{k+2})^m + (q-1) \left[(\lambda_{h,4})^m + (\lambda_{h,5})^m \right]. \quad (21)$$

For the symmetric homeomorphic expansion, $P((Ch)_{(k,k),cyc.,m}, q)$ and $P((Ch)_{(k,k),Mb.,m}, q)$ and the resultant W and $\mathcal{B}$ were given in [23]. Results for P , W , and $\mathcal{B}$ for the cyclic and Möbius $L_y = 2$ triangular strips were given in [24] and P for the cyclic $L_y = 2$ triangular strips also in [27]. Note that for the $L_y = 2$ Möbius triangular strip and homeomorphically expanded square strip in eq. (19) two of the c_j 's are nonpolynomial algebraic functions of q .

We comment on how the nonpolynomial algebraic roots in the various P 's yield polynomials in q . As is evident from (6), for a given strip, the λ_j 's arise as roots of the equation $\mathcal{D} = 0$. In general, $\mathcal{D}$ contains some number of factors of linear, quadratic, cubic, etc. order in x . Consider a generic factor in $\mathcal{D}$ of r 'th degree in x : $(1 + f_1 x + f_2 x^2 + \dots + f_r x^r)$, where the f_j 's are polynomials in x . This yields r λ_ℓ 's as roots of the equation $\xi^r + f_1 \xi^{r-1} + \dots + f_r = 0$. The expressions in P involving these roots are symmetric polynomial functions of the roots, namely terms of the form

$$s_m = \sum_{\ell=1}^r (\lambda_\ell)^m \quad (22)$$

⁶ A homeomorphic expansion of a graph consists in the addition of degree-2 vertices to bonds of that graph.

and, for certain Möbius strips (as well as strips with FBC_x), terms of the form

$$\frac{(\lambda_a)^m - (\lambda_b)^m}{\lambda_a - \lambda_b} = \sum_{k=0}^{m-1} (\lambda_a)^{m-1-k} (\lambda_b)^k. \quad (23)$$

We can then apply a standard theorem (e.g. [32]) which states that a symmetric polynomial function of the roots of an algebraic equation is a polynomial in the coefficients, here f_ℓ , $\ell = 1, \dots, r$; hence this function is a polynomial in q . For example, for s_r , one has the well-known formulas (due to Newton) $s_1 = -f_1$, $s_2 = f_1^2 - 2f_2$, $s_3 = -f_1^3 + 3f_1f_2 - 3f_3$, etc.

Concerning question (i), in general we find that for a given type of lattice strip (or homeomorphic expansion thereof), the terms a_j in (3) are the same for the cyclic and Möbius strips. This can be understood as follows. Recall that the linear equation for the elongation of the strip, together with corresponding equations for related strips produced via deletion-contraction operations, form the matrix equation $Tv_r = v_0$ (eq. (2.29) of Ref. [20]), where v_r is the vector comprised by the generating function Γ for the strip of interest, together with generating functions for the above-mentioned related strips, and v_0 is a vector of chromatic polynomials for beginning forms of these strips. The denominator $\mathcal{D}$ of the generating function Γ is then given as $\mathcal{D} = \det(T)$. The set of generating functions in the vector v_r does depend on the longitudinal boundary conditions, as is clear from the fact that it has larger dimension for periodic than free BC_x 's. However, it is the same for cyclic and Möbius BC_x 's (the elongation is a local operation); the difference here is in the chromatic polynomials in v_0 . For example, in the simplest case of the $L_y = 2$ square strip, the minimal subgraphs, viz., for $m = 2$, are a single square, or equivalently, circuit graph C_4 for PBC_x but the complete graph K_4 for TPBC_x (Möbius).⁷ Hence, $\mathcal{D}$ is the same, while $\mathcal{N}$ is different, for a given strip with cyclic and Möbius longitudinal BC 's. Since the λ_j 's in $\mathcal{D}$ that contribute to P are identical to the a_j 's in eq. (3) (cf. eq. (2.20) in [21]), this enables one to understand that these a_j 's are the same for the cyclic and Möbius strips of a particular type, given the observed property that in the cases where certain λ_j 's do not contribute to P , these are the same for the cyclic and Möbius cases.

The property that the a_j 's are the same for these two cases also answers questions (ii) and (iii). The term a_j that has maximal magnitude in a region of the q plane determines W in that region, and hence the locus $\mathcal{B}$ is the solution of the equation $|a_j| = |a_{j'}|$ of terms of maximal magnitude. It follows that W and $\mathcal{B}$ are the same for the cyclic and Möbius versions of a particular kind of strip. The definition (1) and the equality of the W functions for the cyclic and Möbius strips of a given type G_s imply that, as noted above, as n increases, the h_{n-j} 's for these two strips become equal for progressively larger values of j .

⁷The complete graph K_p is the graph consisting of p vertices all adjacent to each other.

On question (iv), since the c_j are functions of both the λ_j 's in $\mathcal{D}$ and the coefficient functions A_j in $\mathcal{N}$, and since the latter differ for cyclic and Möbius BC_x 's, the c_j 's will also, in general, differ. However, a subset may be the same. In particular, the coefficient c_j of the term which is dominant in region R_1 is always unity, as a consequence of the property that $h_n = 1$ in eq. (2). Hence, *a fortiori*, this coefficient is the same for cyclic and Möbius BC_x 's. In cases where the leading a_j is a nonpolynomial algebraic function of q , the requirement that P be a polynomial implies that certain other coefficients must also be unity. Thus, for example, c_6 and c_7 are both equal to 1 for the $L_y = 3$ square strip, and the $(\lambda_{kg,1})^m$ and $(\lambda_{kg,2})^m$ terms both have coefficient 1 for the kagomé strip.

The situation with the chromatic number, question (v), depends on the particular strip. In the following, we take $L_x = m$ greater than the above-mentioned minimal values where the strips degenerate. The cyclic square strips have $\chi = 2$ for L_x even and $\chi = 3$ for L_x odd, independent of L_y . For Möbius square strips, $\chi = 2$ for $(L_x, L_y) = (e, o)$ or (o, e) , and $\chi = 3$ for $(L_x, L_y) = (e, e)$ or (o, o) , where e and o denote even and odd. Both the cyclic and Möbius kagomé strips above have $\chi = 3$, which shows that the twisting does not necessarily change χ . The cyclic and Möbius homeomorphically expanded square strips [eqs. (21) and (19)] both have $\chi = 3$ for odd k , independent of m , while for even k , the cyclic strip has $\chi = 2, 3$ for m even, odd, and the Möbius strip has $\chi = 2, 3$ for m odd, even, respectively.

For a given type of strip G_s , define $C = \sum_{j=1}^{N_a} c_j$. For the $L_y = 2$ square strips, $C = q(q-1)$ and 0 for the cyclic and Möbius cases [9, 27]. This is also true for the $L_y = 2$ cyclic and Möbius triangular strips, respectively, as also for the cyclic and twisted homeomorphically expanded $L_y = 2$ square strips. For the $L_y = 3$ cyclic square strip, eq. (13) yields $C = q(q-1)^2$, in accord with the generalization $C = P(T_{r=L_y}, q) = q(q-1)^{L_y-1}$ for the cyclic square strip of width L_y . For the $L_y = 3$ square Möbius strip, eq. (7) gives $C = q(q-1)$. The $L_y = 2$ cyclic and Möbius kagomé strips have $C = q^2$ and $C = q$, respectively.

Our comparative study thus elucidates the effect of periodic versus twisted periodic longitudinal boundary conditions on the $T = 0$ Potts antiferromagnet $T = 0$ partition function (chromatic polynomial) for families of lattice strip graphs and homeomorphic expansions thereof. In the future it would also be of interest to extend this study to consider other combinations of longitudinal and transverse boundary conditions.

This research was supported in part by the NSF grant PHY-97-22101.

References

- [1] G. D. Birkhoff, Ann. of Math. **14** (1912) 42.

- [2] R. C. Read, J. Combin. Theory **4** (1968) 52; R. C. Read and W. T. Tutte, “Chromatic Polynomials”, in *Selected Topics in Graph Theory, 3*, (Academic Press, NY, 1988); N. Biggs, *Algebraic Graph Theory* (Cambridge Univ. Press, Cambridge, 1993).
- [3] R. B. Potts, Proc. Camb. Phil. Soc. **48** (1952) 106.
- [4] F. Y. Wu, Rev. Mod. Phys. **54** (1982) 235.
- [5] M. Aizenman and E. H. Lieb, J. Stat. Phys. **24** (1981) 279; Y. Chow and F. Y. Wu, Phys. Rev. **B36** (1987) 285.
- [6] R. Shrock and S.-H. Tsai, Phys. Rev. **E55** (1997) 5165.
- [7] E. H. Lieb, Phys. Rev. **162** (1967) 162.
- [8] J. F. Nagle, J. Combin. Theory **10** (1971) 42; G. A. Baker, Jr., *ibid.* **10** (1971) 217.
- [9] N. L. Biggs, R. M. Damerell, and D. A. Sands, J. Combin. Theory B **12** (1972) 123.
- [10] G. H. Meredith, J. Combin. Theory B **13** (1972) 14.
- [11] N. L. Biggs and G. H. Meredith, J. Combin. Theory B **20** (1976) 5.
- [12] N. L. Biggs, Bull. London Math. Soc. **9** (1977) 54.
- [13] S. Beraha, J. Kahane, and N. Weiss, J. Combin. Theory B **27** (1979) 1; *ibid.* **28** (1980) 52.
- [14] R. C. Read, in Proc. 3rd Caribbean Conf. on Combin. and Computing (1981); Proc. 5th Caribbean Conf. on Combin. and Computing (1988).
- [15] R. J. Baxter, J. Phys. A **20** (1987) 5241; *ibid.* **19** (1986) 2821.
- [16] R. C. Read and G. F. Royle, in *Graph Theory, Combinatorics, and Applications* (Wiley, NY, 1991), vol. 2, p. 1009.
- [17] R. C. Read and E. G. Whitehead, Discrete Math. **204** (1999) 337 and unpublished reports.
- [18] R. Shrock and S.-H. Tsai, Phys. Rev. **E56** (1997) 1342, 3935.
- [19] R. Shrock and S.-H. Tsai, Phys. Rev. **E58** (1998) 4332; cond-mat/9808057.
- [20] M. Roček, R. Shrock, and S.-H. Tsai, Physica **A252** (1998) 505; *ibid.* **A259** (1998) 367.

- [21] R. Shrock and S.-H. Tsai, Physica **A259** (1998) 315.
- [22] R. Shrock and S.-H. Tsai, J. Phys. A **31** (1998) 9641; Physica **A265** (1999) 186.
- [23] R. Shrock and S.-H. Tsai, J. Phys. A Lett. **32** (1999) L195.
- [24] R. Shrock and S.-H. Tsai, Phys. Rev. E, in press; Physica A, in press.
- [25] R. Shrock and S.-H. Tsai, J. Phys. **32** (1999) 5053. (cond-mat/9905431).
- [26] A. Sokal, Combin. Prob. Comput., in press (cond-mat/9904146).
- [27] N. L. Biggs, London School of Economics preprint.
- [28] D. G. Wagner, Waterloo preprint.
- [29] G. Chia, Discrete Math. **172** (1997) 191.
- [30] R. Shrock and S.-H. Tsai, Phys. Rev. **E55** (1997) 6791; *ibid.* **E56** (1997) 2733.
- [31] R. Shrock and S.-H. Tsai, Phys. Rev. **E56** (1997) 4111.
- [32] J. V. Uspensky, *Theory of Equations* (McGraw-Hill, NY 1948), 264.